

## Kingsgate 2002

- Orange Swift  
1/8-8/9. One-nine on most nights, with peaks of 12 on 16/8 and 15 on 18/8.
- Common Swift  
9/5-8/7. One-twenty on most nights, with 22 on 3/6 and 24 on 17/6.
- Six-spot Burnet Moth  
A diurnal species, small numbers amongst the Golf Course rough, usually nectaring on Scabious or Clover flowers, from 20/6 until mid August. .
- December Moth  
One-three on three dates 25/11-16/12.
- Lackey  
Singles on 26/6, 31/7 and 14/8.
- Oak Eggar  
Single females on five dates 21/7-21/8. The males are diurnal and were noted flying by day on several dates from mid July-mid August
- Oak Hook-tip  
One on 26/7.
- Chinese Character  
Singles on nine dates 24/7-25/8.
- Buff Arches  
15/6-11/8. 1-4 on twenty-nine dates.
- Figure of Eighty  
2/6-4/7. 1-3 on seven dates with 4 on 17/6.
- Common Lutestring  
One on 17/7.
- March Moth  
Singles on 16/3, 21/3 and 25/3.
- Rest Harrow  
One on 21/6. The 2<sup>nd</sup> area record.
- Common Emerald  
14/6-9/8. Singles on only five dates in June, then 1-5 on most nights.
- Small Emerald  
Singles on 19/7, 28/7 and 7/8.
- Maiden's Blush  
One on 21/5.
- Blood-vein  
Double brooded. 23/5-17/6, singles on four dates. 26/7-17/8, singles on seven dates.
- Mullein Wave  
Double brooded. 10/5-14/7, 1-2 on seven dates. 30/7-8/9, 1-2 on fifteen dates.

- Small Blood-vein  
16/6-24/8. 1-5 on twenty-five dates, most frequent in July.
- Least Carpet  
19/6-28/8. 1-17 most nights from July with counts of 18-32 on four dates and 45 on 30/7.
- Small fan-footed Wave  
6/7-18/8. 1-3 on sixteen dates.
- Dwarf Cream Wave  
21/6-1/8. 1-3 on sixteen dates.
- Small Dusty Wave  
10/5-20/9. Noted on only two dates in May with 1-6 on seventeen dates in June/July and a peak of 17 on 17/6. 1-6 most nights thereafter with 10 on 29/8 and 8 on 31/8.
- Single-dotted Wave  
15/6-3/9. 1-7 on numerous dates, especially in August. Record counts of 12 on 28/7 and 16 on 30/7.
- Satin Wave  
Double brooded. Singles on 15/6 and 17/6. Five singles from 24/7-18/8 with 3 on 26/7.
- Treble Brown Spot  
8/6-8/8. 1-5 on twenty-one dates with 8 on 17/6.
- Riband Wave  
15/6-19/9. 1-4 most nights in June, 1-12 most nights in July/August with 14-37 on thirteen dates and a record 67 on 30/7. Only two singles in September.
- Vestal  
Immigrant. One on 7/10.
- Gem  
Immigrant. A total of 10 noted in two periods of migrant activity. Five from 8/8-24/8 and five from 14/10-26/10. Singles on six dates with 2 on 8/8 and 26/10.
- Red Twin-spot Carpet  
No 1<sup>st</sup> generation records. 21/7-18/8. 1-2 on seven dates with 4 on 28/7.
- Dark-barred Twin-spot Carpet  
Double brooded. One on 17/5. Then 2 on 28/7 with singles on 29/7, 30/7 and 7/8.
- Silver-ground Carpet  
One on 24/5.
- Garden Carpet  
15/5-6/10. At least double brooded but overlapping. 15/5-6/10. 1-2 on just four dates in May, thereafter 1-12 on most nights.
- Shaded Broad-bar  
6/7-13/8. Singles on six dates with 2 on 28/7.
- Ruddy Carpet  
One on 17/6. The 2<sup>nd</sup> area record.
- Common Carpet  
Double brooded. 12/5-25/6. 1-4 on nine dates with 6 on 17/6.  
7/7-8/9. 1-5 on twenty-nine dates with 7 on 30/7.

Wood Carpet  
One on 18/6. The 3<sup>rd</sup> area record and the first since 1994.

Galium Carpet  
Double brooded. One on 17/6 and then singles on six dates from 15/8-31/8.  
Appears to have recently established itself in this area, first recorded in 1998.

Yellow Shell  
17/6-8/8. Singles on six dates with 2 on 7/7. A poor year.  
A common moth with only a casual attraction to light traps.

Mallow  
29/9-11/11. Singles on six dates with 2 on 26/10.

Shoulder Stripe  
One on 27/4. Only the 3<sup>rd</sup> area record.

Streamer  
One on 3/4. The 2<sup>nd</sup> area record.

Dark Spinach  
Singles on 24/7, 30/7 and 18/8.

Purple Bar  
Double brooded. 21/5-17/6. Singles on five dates. 28/7-28/8. 1-2 on eight dates.

Phoenix  
Singles on 28/7 and 13/8.

Spinach  
17/6-24/7. Singles on five dates

Small Phoenix  
One on 26/7.

Common Marbled Carpet  
Double brooded. 27/5-30/6. 1-5 on thirteen dates with 9 on 17/6 and 8 on 21/6.  
18/8-14/10. 1-3 on thirteen dates.

Barred Yellow  
One on 10/6, 2 on 17/6 and a further single on 7/7.

Blue-bordered Carpet  
One on 31/7.

Grey Pine Carpet  
Double brooded. 2/6-25/6. Singles on five dates  
11/8-6/10. Singles on four dates.

July Highflyer  
One on 26/7.

Small Waved Umber  
One on 13/8.

Fern  
One on 12/8.

Pretty Chalk Carpet  
One on 16/8.

November Moth  
26/10-23/11. 1-2 on seven dates with 4 on 5/11.

Winter Moth  
One on 16/12.

Small Rivulet  
Singles on 22/7 and 24/7.

Barred Rivulet  
One on 18/8. The 2<sup>nd</sup> area record.

Grass Rivulet  
One on 16/6. The 1<sup>st</sup> area record.

Sandy Carpet  
One on 17/6. Surprisingly only the 2<sup>nd</sup> area record.

Maple Pug  
One on 28/7. The 1<sup>st</sup> area record.

Haworth's Pug  
21/6-8/8. Singles on five dates.

Toadflax Pug  
Singles on 16/7, 29/7 and 13/8.

Mottled Pug  
17/5-28/6. 1-5 on twenty-one dates.

Lime-speck Pug  
At least double brooded but overlapping. 17/5-2/10. 1-3 on just seven dates in May, with 1-4 on fourteen dates in June. 1-8 most nights in July/August with 10 on 29/7 and 14 on 30/7. 1-2 on just five dates during September.

Freyer's Pug  
2/6-7/7. 1-2 on eight dates with 7 on 17/6.

Wormwood Pug  
12/7-18/8. 1-4 on twenty-two dates with 6 on 30/7. In addition, a late specimen on 1/9.

Currant Pug  
Double brooded. 23/5-29/6. Singles on four dates with 2 on 29/6.  
22/7-23/8. 1-2 on twelve dates.

Common Pug  
Double brooded. 17/5-25/6. 1-8 on twenty dates with 12 on 17/6.  
One on 16/8 was the only 2<sup>nd</sup> generation record.

White-spotted Pug.  
One on 16/6. Singles on four dates 28/7-16/8.

Grey Pug  
31/5-17/6. Singles on five dates. In addition, one on 22/7.

Tawny Speckled Pug  
30/7-4/9. 1-4 on seventeen dates.

Bordered Pug  
28/7-29/8. Singles on nine dates with 3 on 4/8.

- Shaded Pug  
30/5-7/7. 1-7 on nineteen dates with 19 on 17/6 and 14 on 21/6.  
Late records of 1 on 30/7 and 2 on 31/7.
- Yarrow Pug  
Singles on 21/7, 28/7 and 31/7.
- Plain Pug  
21/7-21/8. Singles on ten dates.
- Oak-tree Pug  
One on 17/5.
- Juniper Pug  
One on 24/6. The 2<sup>nd</sup> area record.
- Cypress Pug  
One on 17/5 is by far the earliest area record. 25/7-19/9. 1-8 most nights with 16 on 16/8,  
20 on 18/8 and 14 on 23/8.
- V-Pug  
Singles on 25/7 and 11/8.
- Sloe Pug  
30/5-16/6. Singles on four dates with 2 on 6/6.
- Green Pug  
30/5-7/7. 1-9 on seventeen dates with 12 on 16/6 and 15 on 17/6.
- Double-striped Pug  
Double brooded. 16/3-11/5. 1-5 on eight dates. One on 24/6.  
8/7-23/9. 1-4 on twenty-eight dates with 7 on 11/8.
- Treble-bar  
Double brooded. Singles on three dates 19/5-21/6 and on four dates 24/7-30/8.
- Small Yellow Wave  
One on 17/6.
- Yellow-barred Brindle  
Double brooded. 20/5-3/7. 1-2 on nine dates.  
29/7-7/9. Singles on eleven dates with 2 on 18/8.
- Magpie  
2/6-28/8. 1-6 on most nights until 26/6 with 9 on 17/6, then none until 12/7. 1-6 on eight  
dates in July and 1-9 on most nights in August.
- Clouded Border  
Singles on 30/7 and 10/8.
- Scorched Carpet  
One on 18/8.
- Peacock  
Singles on 8/8 and 18/8.
- Sharp-angled Peacock  
Double brooded. One on 17/6. Singles on 28/7, 30/7, 3/8 and 17/8.

Tawny Barred Angle

One on 4/7.

Latticed Heath

Singles on 21/6, 24/7 and 8/8.

Brimstone

At least double brooded. 27/4-25/6. 1-8 most nights with 10 on 17/6.  
7/7-13/9. 1-7 most nights in July and 1-15 on most nights during August with 18 on 7/8  
and 21 on 18/8. Only noted as 1-2's during September.

Large Thorn

One on 8/9.

Canary-shouldered Thorn

One on 1/9.

Early Thorn

Only 2<sup>nd</sup> generation examples recorded. Singles on seven dates 16/7-6/8.

Scalloped Oak

21/6-17/8. Only two singles in June, then 1-6 on most nights.

Swallow-tailed Moth

One on 26/6 and then 1-7 on most nights 12/7-4/8.

Feathered Thorn

4/11-11/11. Singles on three dates with 3 on 5/11.

Peppered

9/6-13/8. 1-3 on sixteen dates with 4 on 17/6.

Dotted Border

16/3-3/4. Singles on five dates.

Waved Umber

12/5-24/6. 1-2 on ten dates.

Willow Beauty

Five singles 15/6-25/6 is unusual. 24/7-29/9. 1-10 on most nights during July/August,  
with 14-26 from 14/8-17/8 and 33 on 18/8. 1-2 on nine dates during September.

Mottled Beauty

15/6-14/7. 1-3 on twelve dates.

Engrailed

One on 3/4.

Bordered White

One on 17/6.

Common White Wave

One on 17/6.

Clouded Silver

2/6-12/8. 1-5 on twelve dates.

Light Emerald

Double brooded. 15/5-26/6. Singles on 15/5 and 23/5 then 1-3 most nights in June, with 5 on 5/6.  
7/8-23/9. 1-5 most nights during August and 1-2 on just nine dates during September.

Yellow Belle.

Double brooded. Singles on 28/5 and 31/5. Further singles on 23/8, 28/8 and 31/8.

Convolvulus Hawk-moth

A Mediterranean immigrant. One on 13/8.

Privet Hawk-moth

One on 15/6.

Lime Hawk-moth

21/5-17/7. Singles on nine dates. A record year.

Eyed Hawk-moth

21/5-8/8. 1-2 on eighteen dates, totalling 21. Also a record year.

Poplar Hawk-moth

21/5-18/8. 1-4 on nineteen dates, totalling 25 is slightly below average.

Hummingbird Hawk-Moth

A Mediterranean immigrant.

Only noted by day feeding on cliff-top flowers. Singles on 15/6, 28/9 and 10/10.

Striped Hawk-moth

A Mediterranean immigrant. Different singles on 15/6, 17/6 and 19/6. The 2<sup>nd</sup>-4<sup>th</sup> records.

Elephant Hawk-moth

Singles on 16/6 and 7/7.

Small Elephant Hawk-moth

2/6-18/6. Singles on just four dates with 2 on 15/6.  
A total of 109 were recorded in 1998 and 99 in 1999!

Buff-tip

Singles on 17/5, 15/6 and 7/7.

Puss Moth

One on 15/6.

Pebble Prominent

Double brooded. Singles on 7/6 and 17/8.

Swallow Prominent

Double brooded. 27/4-10/6. Singles on five dates.  
30/7-24/8. Singles on five dates with 4 on 7/8 and 2 on 8/8.

Coxcomb Prominent

Singles on 12/7 and 18/8.

Maple Prominent

One on 15/8.

Pale Prominent

Double brooded. Singles on 16/6 and 18/8.

Vapourer  
Singles on 17/8 and 8/9. Commonly seen flying by day, only a casual attraction to light.

Pale Tussock  
Singles on 17/5, 26/5, 30/5 and 2/6. Rather surprisingly a record year.

Brown-tail  
14/7-2/8. 1-3 on most nights.

Yellow-tail  
24/7-3/8. Singles on seven dates with 2 on 27/7.

Dingy Footman  
20/7-4/9. 1-4 on eighteen dates.

Scarce Footman  
14/7-19/8. 1-9 on most nights.

Buff Footman  
One on 7/8. The 1<sup>st</sup> record for the area.

Common Footman  
21/6-7/8. After just three singles in June, noted on every night, usually 5-19.

Garden Tiger  
One on 28/6 and then 1-3 on eleven dates from 26/7-8/8.

White Ermine  
15/5-24/6. Singles on seven dates.

Buff Ermine  
14/5-6/8. 1-2 on five dates in May, and then 1-9 on most nights with 18 on 17/6 and 12 on 21/6.

Muslin Moth  
3/4-17/5. Singles on five dates with 2 on 15/5. Only the males attracted to light.

Ruby Tiger  
Only noted in the 2<sup>nd</sup> generation. 8/7-2/8. 1-4 on eleven dates.

Cinnabar  
19/5-13/7. Two on 19/5 and then singles on five dates.

Kent Black Arches  
Singles on 21/6 and 27/7 is exceptionally poor.

Short-cloaked Moth  
Singles on 15/6 and 12/7 is also very poor.

Least Black Arches  
Just a single on 13/5. Very poor. The 2002 weather was presumably unsuitable for the three Nola species!

White-line Dart  
Singles on 2/8, 7/8 and 10/8 with 2 on 13/8 is by far the worst year on record.

Garden Dart  
Continuing it's marked decline over the last few years. One on 14/8.

- Turnip  
17/6-10/9. Singles on just nineteen dates with 4 on 28/7. Further singles on 13/10 and 10/11.
- Dark Sword-grass  
Immigrant. 12/5-5/11. 1-4 on forty dates totalling just 60. Most frequent during July/August (39). The worst year since 1997.
- Heart and Club  
10/6-28/7. 1-18 on most nights with a peak of 24 on 4/7.
- Heart and Dart  
6/5-22/8. A poor year. Every night during June/July, including 40-99 on twenty-three dates with 112 on 21/6 and 103 on 30/6. Only in single figures from 18/7.
- Shuttle-shaped Dart  
At least double brooded. 27/4-29/6. 1-9 on most nights.  
17/7-5/10. 1-15 on most nights during July/August with 21 on 8/8. 1-3 on eight dates in September and on three dates in October.
- Sand Dart  
Singles on 22/6 and 27/6 with 2 on 17/7.
- Flame  
14/6-17/7. 1-3 on twenty-one dates with 4 on 15/6. One of the occasional 2<sup>nd</sup> brood on 10/9.
- Flame Shoulder  
Double brooded. 9/5-6/7. 1-2 on fourteen dates. 17/7-1/9. 1-8 on most nights.
- Large Yellow Underwing  
Very common. One on 20/3 is the earliest record for the area. 17/5-10/11. A total of 3454, represents a poor year (5316 in 2001) or a good year, depending on your view – mine is definitely the latter! 1-19 on most nights during June with 148 on 30/6. Recorded on every night from July-September, typically 15-72 with peaks of 204 on 5/9, 600 on 6/9 and 268 on 7/9. Although regular regular throughout October, a maximum of only 12.
- Lesser Yellow Underwing  
22/6-11/11. Thankfully an extremely poor year. 1-24 on most nights during July with 33 on 26/7. In August, 1-4 until 13/8 then none until 24/8. 1-28 on most nights thereafter.
- Broad-bordered Yellow Underwing  
4/7-1/10. 1-2 on sixteen dates.
- Lesser Broad-bordered Yellow Underwing  
18/7-17/9. 1-11 on most nights during July and September. Recorded on every night during August, including 20-74 on fourteen dates from 7/8-21/8, with a peak of 97 on 17/8.
- Least Yellow Underwing  
5/7-18/9. 1-5 on thirty-two dates.
- Double Dart  
One on 4/7. The 1<sup>st</sup> area record.
- Pearly Underwing  
Immigrant. 21/5-24/11. 1-2 on twenty dates, fairly evenly distributed throughout the period, totalling just 27.
- Ingrailed Clay  
One on 17/6. The 2<sup>nd</sup> record.

Small Square-spot

Double brooded. 21/5-26/6. 1-4 on most nights from 6/6.  
21/7-18/9. 1-3 on just fourteen dates.

Setaceous Hebrew Character

Double brooded. 19/5-2/7. 1-4 on many dates.  
31/7-2/10. 1-9 on most nights with 11-15 on five dates and a peak of 19 on 18/8.

Double Square-spot

17/6-29/7. Singles on seven dates.

Six-striped Rustic

Two on 4/8 with singles on 14/8 and 17/8.

Square-spot Rustic

13/7-14/10. Most nights (every night 1/8-13/9) with 15-37 on twenty dates and 47 on 18/8.  
Only in single figures from 10/9. A poor year.

Nutmeg

Singles on 8/6, 15/6 and 30/6. Then 1-2 on fifteen dates 17/7-24/8.

Shears

27/5-5/7. Singles on eight dates. A record year.

White Colon

Singles on 31/5 and 8/7.

Cabbage

Double brooded. 27/4-22/6. Singles on twelve dates. 1/7-16/9. 1-2 on ten dates in July with 5  
on 31/7, after which 1-8 on most nights with 10 on 11/8.

Dot

17/6-13/8. Recorded on most nights with counts of 10-17 on eight dates during July.

Light Brocade

Singles on 28/5, 6/6 and 15/6.

Pale-shouldered Brocade

One on 4/6.

Bright-line Brown-eye

1/5-23/9. 1-2 on most nights in May and 1-10 on most nights during June. Noted on every  
night in July with 10-19 on fourteen dates. 1-13 on most nights in August and 1-4 on most  
nights during September.

Dog's Tooth

15/6-18/8. Singles on six dates.

Broad-barred White

16/6-21/8. Singles on just five dates. A much decreased species.

Campion

Double brooded. 21/5-15/6. 1-2 on five dates.  
17/7-5/9. 1-2 on seventeen dates with 4 on 24/7.

Varied Coronet

Singles on 21/6, 30/6 and 13/7.

- Lychnis  
Double brooded. 21/5-7/7. Singles on six dates with 2 on 14/6.  
24/7-8/9. Singles on eight dates with 2 on 8/9.
- Hedge Rustic  
One on 5/9.
- Pine Beauty  
Singles on 14/5 and 17/5.
- Small Quaker  
One on 3/4. By far the worst year on record and an indication of the poor spring weather.
- Common Quaker  
3/3-31/5. 1-9 on many nights. Another common species finding recent springs unfavourable.
- Clouded Drab  
21/3-10/5. Singles on only six dates with 2 on 4/4.
- Hebrew Character  
20/3-20/5. 1-6 on many nights throughout with a peak of 10 on 3/4.
- White-point  
Immigrant. Probably temporarily established, at least in the Pegwell area.  
One on 25/5, 3 on 8/6 and 1 on 23/6 are for here unusual spring records.  
More typically, singles on five dates from 5/8-18/8 with 2 on 26/8 and 4 on 2/9.  
Finally, 1 on 7/10. A yearly total of 17 is slightly below the average of recent years.
- Brown-line Bright-eye  
6/7-18/8. 1-2 on eleven dates.
- Clay  
6/7-14/8. 1-6 on most nights.
- Delicate  
Immigrant. Singles on 7/7 (the 1<sup>st</sup> July record) and 14/10.
- Southern Wainscot  
One on 4/7.
- Smoky Wainscot  
15/6-14/8. 1-3 on five dates in June and on every night during July, with 10-20 on thirteen dates and a peak of 37 on 27/7. 1-5 on most nights during August.
- Common Wainscot  
Double brooded. 19/5-15/7. 1-4 on fourteen dates with 9 on 16/6.  
14/8-9/9. 1-5 on most nights. Also an isolated individual on 5/10. Poor.
- L-Album Wainscot  
Established on Thanet since the mid 90's. Double brooded. 18/6-30/7. 1-4 on many nights.  
28/8-28/10. 1-7 on most nights with 12 on 19/9.
- White-speck  
Immigrant. Sporadic in its appearance. Singles on 6/10, 14/10, and 31/10 with 2 on 26/10.  
A further single on 1/12.
- Wormwood  
One on 22/7.

- Chamomile Shark  
Singles on 1/5, 10/5 and 25/5.
- Shark  
7/6-7/7. Singles on eight dates with 4 on 2/7 and 2 on 7/7.
- Star-wort  
One on 7/7 and then further singles on four dates 10/8-24/8.
- Deep-brown Dart  
One on 4/10. Exceptionally poor, normally a common autumn moth here.
- Black Rustic  
Another colonist of the mid 90's. 25/9-2/10. Singles on six dates with 3 on 1/10 and 2 on 2/10.
- Blair's Shoulder-knot  
18/9-10/11. Singles on just three dates during September, thereafter, 1-6 on most nights.
- Feathered Brindle  
One on 28/9.
- Pale Pinion  
One on 18/9. The 1<sup>st</sup> area record.
- Early Grey  
5/3-25/5. Most nights, typically 1-8 with 9-18 on six dates.
- Green-brindled Crescent  
26/10-10/11. Singles on just four dates.
- Feathered Ranunculus  
21/8-28/10. Noted on three dates during August and then 1-16 on most nights. As with most of the autumnal species, a poor year.
- Satellite  
Singles on 29/10, 4/11 and 30/11.
- Chestnut  
One on 30/3. Two on 11/10 and then singles on 30/10, 5/11 and 3/12.
- Brick  
Singles on 16/9 and 3/10. Then 1-2 on fourteen dates 26/10-3/12.
- Yellow-line Quaker  
One on 24/11.
- Beaded Chestnut  
Singles on 17/9 and 6/10 is exceptionally poor.
- Centre-barred Sallow  
One on 6/9.
- Lunar Underwing  
16/9-7/10. 1-8 on most nights. Very poor.
- Poplar Grey  
15/6-7/8. Singles on ten dates with 2 on 14/7.

Sycamore  
17/6-4/8. Singles on seven dates.

Grey Dagger  
17/6-11/8. 1-3 on eleven dates with 6 on 30/7.

Knot Grass  
Singles on 30/7, 11/8 and 24/8.

Tree-lichen Beauty  
A rare immigrant. Singles on 12/8 and 15/8. The 2<sup>nd</sup> and 3<sup>rd</sup> area records.

Marbled Beauty  
15/7-7/9. 1-11 on most nights with a peak of 16 on 19/8.

Marbled Green  
20/7-30/8. 1-8 on most nights.

Copper Underwing  
6/8-6/10. 1-6 on most nights.

Svensson's Copper Underwing  
One on 18/8. The 2<sup>nd</sup> area record.

Mouse  
22/7-25/9. 1-8 on most nights, commonest during August.

Old Lady  
Singles on 24/8, 25/8 and 28/8. A poor year.

Brown Rustic  
One on 23/6.

Straw Underwing  
30/7-7/9. 1-13 on most nights with peaks of 19 on 17/8 and 24 on 18/8.

Small Angle Shades  
Singles on 20/6 and 15/7, are the first to be recorded at Kingsgate since 1990!

Angle Shades  
Resident and supplemented by immigration, particularly during the autumn.  
One on 25/3. 27/4-26/6. 1-5 most nights during April/May and 1-10 most nights during June.  
Singles on 5/7 and 23/7 were the only July records.  
6/8-16/12. 1-6 on most nights with 25 on 6/9, 14 on 7/9 and 11 on 26/10. A poor year.

Dingy Shears  
One on 8/7.

Lesser Spotted Pinion  
One on 16/8.

Dun-bar  
16/7-14/8. 1-3 on eighteen dates.

Dark Arches  
27/5-20/8. Every night from 13/6. 1-30, most frequent in July, with a peak of 43 on 15/7.

Light Arches  
14/6-24/7. Singles on just four dates. A very marked decline in recent years.

Reddish Light Arches  
One on 26/6.

Crescent Striped  
One on 8/7.

Clouded Brindle  
14/6-27/6. Singles on four dates.

Dusky Brocade  
8/6-30/7. 1-6 on many nights with a peak of 9 on 8/7.

Rustic Shoulder-knot  
18/5-25/6. Singles on six dates with 2 on 4/6.

Slender Brindle  
One on 8/7.

Marbled Minor  
10/5-1/7. 1-15 on most nights in May and 1-30 on most nights in June with of 35 on 17/6.

Tawny Marbled Minor  
19/5-16/7. 1-19 on most nights with peaks of 28 on 17/6 and 26 on 21/6.

Middle-barred Minor  
Singles on 17/6 and 21/6.

Cloaked Minor  
17/7-4/9. 1-6 on seven dates during July, 1-10 on most nights thereafter.

Rosy Minor  
Singles on 24/7, 30/7 and 6/8.

Common Rustic  
8/7-12/9. 1-8 on every night during July. Recorded on every night during August, with 21-72 on thirteen dates and a peak of 83 on 17/8. 1-2 on three dates in September. Poor.

Lesser Common Rustic  
19/7-19/8. Obvious singles on five dates with 2 on 8/8.

Dusky Sallow  
16/7-14/8. 1-2 on ten dates with 4 on 27/7.

Flounced Rustic  
19/7-4/10. Most nights, typically 10-36. Poor.

Sandhill Rustic - demuthi  
One on 13/8. The 1<sup>st</sup> area record. A very local Salt-marsh species, the nearest colony is at Faversham Creek

Saltern Ear  
One on 15/8.

Rosy Rustic  
11/8-18/9. Singles on just five dates.

Crescent  
One on 19/8.

Bulrush Wainscot

One on 21/8. The 4<sup>th</sup> record of this marshland species.

Brown-veined Wainscot

One on 12/8.

Webb's Wainscot

One on 23/8. The 4<sup>th</sup> record of this marshland species.

Large Wainscot

One on 5/11. The latest record for the area.

Fen Wainscot

One on 28/7.

Treble Lines

15/5-23/6. 1-14 on most nights with a peak of 16 on 31/5.

Uncertain

15/6-17/8. 1-7 on most nights in June and August. 3-20 every night in July with 26 on 8/7.

Rustic

4/7-18/8. 1-5 most nights in July with 1-2 most nights during August.

Small Mottled Willow

Immigrant. One on 17/6. The 11<sup>th</sup> record for the area.

Vine's Rustic

Double brooded. 17/5-28/6. 1-4 most nights in May and 1-10 on most nights in June. Singles on three dates in July. 3/8-29/9. 1-21 on most nights, single figures only from 8/9.

Mottled Rustic

28/5-26/7. 1-20 on most nights with 27 on 20/6 and 29 on 24/6.

Pale Mottled Willow

An isolated single on 4/4. Then noted from 6/5-29/10. 1-4 on most nights from May-July and 1-6 on most nights during August/September. 1-3 on just six dates in October.

Bordered Sallow

15/6-27/7. Singles on just four dates. Very poor.

Scarce Bordered Straw

Immigrant. One on 17/6 is the 1<sup>st</sup> spring record. More typically, singles on 10/9, 16/9, 14/10, 26/10 and 5/11. A total of six is an average showing of this formally scarce migrant.

Bordered Straw

Immigrant. One on 16/6, 7 on 17/6, 2 on 19/6, 1 on 30/6 and 2 on 18/8. A total of thirteen is a record. Typically 2-5 a year although none in 2001 and 11 in 2000.

Cream-bordered Green Pea

One on 30/7.

Oak Nycteoline

One on 31/3.

Golden Twin-spot

A rare immigrant. Singles on 30/7 and 30/9. The 9<sup>th</sup> and 10<sup>th</sup> area record. Something of a Thanet speciality in recent years, most of the UK records occurring here.

- Ni Moth  
An uncommon immigrant. One on 26/8. The 2<sup>nd</sup> area record.
- Burnished Brass  
Double brooded. 14/6-30/7. 1-4 on twenty-two dates with 5 on 8/7.  
16/8-25/9. 1-2 on thirteen dates with 4 on 29/8.
- Gold Spot  
One on 24/8.
- Silver Y  
A common immigrant. A total of 479 from 11/5-3/12. Average numbers May-August after which very poor. 1-4 on sixteen dates in May with 14 on 21/5. 1-17 on most nights from June-August with peaks of 26 on 13/6, 40 on 17/6, 22 on 17/8 and 27 on 18/8.  
Only noted in single figures from September.
- Dark Spectacle  
Only three records up to 1999, now common. Double brooded, the broods overlapped.  
26/5-8/9. 1-4 on many dates throughout with 7 on 8/7.
- Spectacle  
Again double brooded with no clear differentiation between broods.  
20/5-5/9. 1-5 on many dates throughout.
- Red Underwing  
Different singles on 30/9 and 1/10.
- Mother Shipton  
A day flying species that can be found with a little effort on the golf course.  
19/5-13/6. 1-3 on seven dates.
- Herald  
Singles following winter hibernation on 15/3 and 25/4.  
Summer brood specimens from 3/7-31/8. Singles on six dates with 2 on 17/7.
- Straw Dot  
15/6-14/8. Singles on four dates with 2 on 7/8.
- Snout  
Double brooded. 15/6-30/7. Singles on eight dates. 14/8-28/9. Six singles with 2 on 7/9.
- Fan-foot  
26/6-28/8. Singles on six dates with 2 on 13/7 and 4 on 27/7.
- Small Fan-foot  
One on 17/7.
- Dotted Fan-foot  
One on 16/7. The 1st area record.

## Microlepidoptera

Strict family Order has not been followed, as most people are likely to have some interest in the Pyralids (the only micros that I comprehensively recorded) and to a lesser extent the Tortrix. Family Order resumes after these.

### Pyralids

*Chrysoteuchia culmella* – Garden Grass-veneer

22/5-18/8. Common, every night from 13/6-10/8.

*Crambus pascuella*

15/6-9/8. 1-6 on many nights, most frequent during July.

*Crambus lathoniellus*

15/5-4/7. 1-10 on fourteen dates.

*Crambus perlella*

19/6-12/9. Common, most nights from 4/7.

*Agriphila straminella*

7/7-28/8. Small numbers on most nights.

*Agriphila tristella*

27/7-9/9. Common, most nights.

*Agriphila latistria*

Singles on 17/8, 18/8 and 29/8 with 2 on 3/9.

*Agriphila geniculea*

27/7-23/9. Common, most nights.

*Catoptria pinella*

Singles on 28/7 and 12/8.

*Catoptria falsella*

Singles on 28/7 and 6/8.

*Pediasia contaminella*

One on 20/8. The 3<sup>rd</sup> area record.

*Platytes alpinella*

Singles on 29/7 and 13/8.

*Platytes cerussella*

Two on 15/6. Recorded on single dates in the last three years.

*Acentria ephemerella* – Water Veneer

A marshland species, the males of which occasionally swarm far from water.

A record 78 on 24/7 with 23 on 25/7. Then 1-5 noted on eight dates from 27/7-19/8.

*Scoparia subfusca*

15/6-16/7. 1-4 on six dates.

*Scoparia pyralella*

5/6-12/7. 1-3 on fourteen dates with 7 on 17/6.

*Scoparia ambigualis*

5/6-15/8. 1-11 on most nights.

- Eudonia pallida*  
One on 14/6.
- Dipleurina lacustrata*  
17/6-28/7. 1-5 frequently recorded.
- Eudonia angustea*  
Double brooded. 21/3-5/7 and 9/8-5/11. Common, most nights.
- Eudonia mercurella*  
17/6-18/8. Small numbers most nights from July.
- Evergestis forficalis* – Garden Pebble  
Double brooded. 6/5-2/7. 1-4 on many nights. 31/7-16/9. 1-10 on most nights.
- Evergestis extimalis*  
15/7-20/9. 1-9 most nights during July/August with peaks of 11 on 28/7 and 17 on 30/7.
- Pyrausta aurata*  
Two on 30/7 and then singles on four dates from 7/8-29/8.
- Ostrinia nubilalis* – European Corn-borer  
15/6-29/7. 1-2 on ten dates.
- Eurrhpara hortulata* – Small Magpie  
31/5-29/8. 1-8 on most nights until the end of July. Only five singles during August.
- Phlyctaenia coronata*  
24/6-7/9. Common, most nights.
- Udea prunalis*  
Singles on 25/6 and 16/7.
- Udea olivalis*  
5/6-5/7. 1-7 on most nights.
- Udea ferrugalis* – Rusty-dot Pearl  
Immigrant. One on 9/6. Then 1-3 on twenty-one dates from 10/8-2/12, totalling 31.
- Nomophila noctuella* – Rush Veneer  
Immigrant. One on 22/5 was the 1<sup>st</sup> May record for the area. Two on 17/6 and 1 on 19/6.  
Noted from 23/7-11/11. 1-6 on forty-six dates with 8 on 8/8. Most frequent during August.  
A year total of 107.
- Pleuroptya ruralis* – Mother of Pearl  
16/7-24/8. 1-9 on most nights.
- Hypsopygia costalis* – Gold Triangle  
8/7-12/8. 1-3 on ten dates with 7 on 27/7.
- Synaphe punctalis* – Slender Dune  
Singles on 16/7 and 24/7. The 3<sup>rd</sup> and 4<sup>th</sup> area records.
- Orthopygia glaucinalis*  
Singles on 21/7 and 29/7 with 2 on 3/8.
- Pyralis farinalis* – Meal Moth  
21/6-8/8. Singles on five dates with 2 on 29/7.

- Aglossa pingualis* – Large Tabby  
14/6-19/8. Singles on four dates.
- Endotricha flammealis* – Double-barred Flame  
6/7-24/8. Common, most nights.
- Galleria mellonella* – Wax Moth  
Singles on 25/7 and 24/8.
- Aphomia sociella* – Bee Moth  
2/6-10/8. 1-8 on most nights. In addition, 1 on 3/9.
- Acrobasis repandana*  
26/6-19/8. Singles on four dates.
- Trachycera suavella*  
27/7-28/8. Singles on eleven dates with 2 on 15/8.
- Trachycera advenella*  
17/7-7/9. 1-8 on most nights.
- Nephopterix angustella*  
One on 5/8. The 3<sup>rd</sup> area record.
- Phycita roborella*  
27/7-31/8. Singles on five dates with 2 on 18/8.
- Dioryctria abietella*  
One on 23/6.
- Hypochoalcia ahenella*  
25/5-17/7. 1-3 on ten dates.
- Myelois circumvoluta* – Thistle Ermine  
15/6-8/8. 1-2 on nineteen dates.
- Euzophera cinerosella*  
One on 8/8. The 3<sup>rd</sup> area record.
- Euzophera pinguis*  
24/7-15/8. Singles on five dates.
- Ephestia parasitella*  
A difficult group of moths, most, if not all appeared to be this species.  
5/6-10/8. Most nights, up to 20.
- Vitula biviella*  
One on 8/8. This and one at Gravesend (also this year) are the only British records away from the Dungeness area.
- Homoeosoma sinuella*  
15/6-31/7. 1-5 on nineteen dates. In addition, 1 on 5/10.
- Phycitodes binaevella*  
25/6-30/7. Singles on seven dates.

## **Tortrix**

*Cochylimorpha alternana*

Singles on 2/8, 10/8 and 17/8. Very local, restricted in the UK to a few East Kent sites.

*Cochylimorpha straminea*

One on 30/5. Further singles on 21/8 and 3/9.

*Agapeta hamana*

9/6-8/8. 1-4 on eighteen dates, most frequent during July.

*Aethes tesserana*

Singles on 14/6, 19/6 and 12/7.

*Aethes hartmanniana*

A scarce species, only seen by day on the golf course and there restricted to one area of rough. Small numbers noted from 23/7 until early August.

*Aethes cnicana*

One on 12/7.

*Aethes rubigana*

One on 15/7 and 2 on 19/7.

*Aethes smeathmanniana*

Singles on 27/5, 17/6 and 27/7.

*Aethes beatricella*

15/6-28/6. 1-6 on most nights.

*Eupoecilia angustana*.

15/6-30/6. 1-2 on five dates. 28/7-19/8. 1-6 on most nights.

*Eupoecilia ambiguella*

Singles on 30/7 and 8/8.

*Cochylis dubitana*

16/7-19/8. 1-3 on eleven dates.

*Cochylis hybridella*

13/6-3/9. Small numbers throughout.

*Cochylis atricapitana*

15/5-19/6. 1-3 on ten dates. 28/7-5/9. 1-3 regularly recorded.

*Pandemis corylana* – Chequered Fruit-tree Tortrix

Singles on 23/6 and 25/7. Not fully recorded.

*Pandemis cerasana* – Barred Fruit-tree Tortrix

15/6-18/8. Occasional in small numbers throughout.

*Pandemis heparana* – Dark Fruit-tree Tortrix

23/6-29/8. Most frequent from mid July. Regularly 1-8.

*Archips podana* – Large Fruit-tree Tortrix

17/6-8/8. Small numbers throughout.

*Cacoecimorpha pronubana* – Carnation Tortrix

14/5-28/9. Singles on ten dates.

*Aphelia paleana* – Timothy Tortrix  
17/6-9/8. Singles on four dates with 3 on 24/7.

*Clepsia spectrana* – Cyclamen Tortrix  
29/7-23/8. Singles on four dates with 2 on 30/7.

*Clepsia consimilana*  
14/6-19/8. 1-9 on most nights.

*Epiphyas postvittana* – Light Brown Apple Moth  
3/3-5/12. Most nights throughout, very common during August/September.

*Adoxophyes orana* – Summer Fruit Tortrix  
Singles on 20/6, 15/7 and 22/7.

*Lozotaeniodes formosanus*  
21/6-18/8. 1-3 on twenty-two dates.

*Lozotaenia forsterana*  
6/6-20/6. Singles on four dates with 3 on 14/6.

*Ditula angustiorana* – Red-barred Tortrix  
28/6-26/7. Frequently 1-5.

*Pseudargyrotoza conwagana*  
14/6-31/8. 1-3 on ten dates.

*Cnephasia stephensiana* – Grey Tortrix  
30/6-28/7. Frequent in small numbers.

*Cnephasia asseclana* – Flax Tortrix  
Occasional singles 21/6-15/8.

*Cnephasia incertana* – Light Grey Tortrix  
17/6-17/7. Occasional in small numbers.

*Aleimma loeflingiana*  
One on 26/6.

*Tortrix viridana* – Green Oak Tortrix  
17/6-26/6. 1-8 on most nights.

*Acleris forsskaleana*  
16/7-21/8. Common throughout.

*Acleris holmiana*  
One on 15/7.

*Acleris sparsana*  
Over-winters. One on 21/3. 2/10-18/11. 1-3 on nine dates.

*Acleris rhombana* – Rhomboid Tortrix  
Singles on 15/8, 27/9 and 2/10.

*Acleris aspersana*  
3/7-22/8. 1-3 on ten dates.

*Acleris notana*  
One on 27/11.

*Acleris variegana* – Garden Rose Tortrix  
19/6-14/10. Common from July.

*Acleris cristana*  
Over-winters. Singles on 25/3 and 10/11.

*Celypha striana*  
6/6-22/8. 1-2 on eight dates.

*Celypha lacunana*  
25/5-3/9. Common throughout.

*Hedya pruniana* – Plum Tortrix  
19/5-7/7. Common.

*Hedya dimidoalba* – Marbled Orchard Tortrix  
24/5-19/6. Occasional – small numbers.

*Hedya salicella*  
One on 31/7.

*Lobesia abscisana*  
12/7-31/7. 1-3 on five dates.

*Lobesia littoralis*  
One on 1/7.

*Eudemis profundana*  
24/7-30/7. Singles on three dates.

*Ancylis achatana*  
24/6-27/7. 1-4 regularly recorded.

*Epinothia abbreviana*  
One on 27/6.

*Epinothia trigonella*  
One on 3/10.

*Crociosema plebejana*  
5/8-10/11. Singles on six dates.

*Gypsonoma dealbana*  
21/6-22/7. Frequent in small numbers.

*Epiblema cynosbatella*  
15/5-21/6. Regularly 1-4.

*Epiblema uddmanniana* – Bramble Shoot Moth  
15/6-16/8. 1-3 on ten dates.

*Epiblema roborana*  
One on 27/7.

*Epiblema foenella*  
21/7-11/8. Singles on six dates.

*Eucosma campoliliana*  
6/7-17/8. 1-5 on most nights.

*Eucosma hohenwartiana*  
21/6-23/7. Occasionally 1-3.

*Eucosma cana*  
17/6-3/8. 1-4 on most nights.

*Spilonota ocellana* – Bud Moth  
5/6-30/7. 1-4 on most nights.

*Pammene aurantiana*  
One on 9/8.

*Pammene fasciana*  
21/6-25/7. Singles on five dates with 3 on 24/7.

*Cydia nigricana* – Pea Moth  
One on 23/7.

*Cydia splendana*  
One on 28/6 and then 1-8 on most nights from 25/7-24/8.

*Cydia pomonella* – Codling Moth  
14/6-12/8. 1-2 on eight dates with 4 on 31/7.

*Dichrorampha alpinana*  
Singles on 27/7 and 1/8.

*Dichrorampha gueneana*  
Recorded by day, from mid-June, on the Golf Course amongst Ox-eye Daisy and Knapweed.

*Dichrorampha plumbana*  
Recorded by day, from mid-June, on the Golf Course amongst Ox-eye Daisy and Knapweed.

### **Other Families**

*Adela reaumurella*  
A day flying species usually in swarms of 20-100, found in bushy areas on the golf course from late April-early June.

*Nemapogon cloacella* – Skin Moth  
Singles on 17/5 and 22/5.

*Monopis weaverella*  
One on 17/5.

*Monopis laevigella* – Skin Moth  
8/6-15/8. Singles on four dates.

*Monopis obviella*  
6/5-11/11. 1-2 occasionally recorded throughout.

*Niditinea fuscella* – Brown-dotted Clothes Moth  
10/5-25/6. Singles on four dates with 2 on 19/5.

*Tinea semifulvella*  
31/5-28/7. Singles on four dates.

*Tinea trinotella*  
8/6-24/7. Singles on five dates with 2 on 25/6.

*Leucoptera laburnella* – Laburnum Leaf Miner

Singles on 25/6, 8/7 and 27/7. Can also be found by beating Laburnum.

*Lyonetia clerkella* – Apple Leaf Miner

Five on 17/6 with singles on 21/6 and 25/6. Then 25/7-1/9. Frequently 1-4 with 8 on 28/7.

*Bucculatrix nigricomella*

One on 24/7.

*Caloptilia betulicola*

One on 18/9.

*Caloptilia rufipennella*

One on 28/3. Singles on four dates 19/7-29/7 with 3 on 27/7.

*Caloptilia stigmatella*

One on 1/4.

*Caloptilia syringella*

Double brooded. Singles on 27/4, 11/5 and 19/5. 14/8-3/9. Singles on six dates.

*Aspilapteryx tringipennella*

One on 27/7 and 2 on 28/7.

*Parornix anglicella*

Singles on 8/8 and 19/8.

*Acrocercops brongniardella*

The caterpillars mine the leaves of Holm Oak, leaving a distinctive silvery looking blister over most of the upper leaf surface. The only moths were as a result of inhabited leaves collected in August, 2 moths emerged on 15/9. A few colonies on the Golf Course.

*Phyllonorycter messaniella*

Singles on 18/5 and 26/7.

*Phyllonorycter geniculella*

Mines Sycamore leaves. One swept from Sycamore on 24/5.

*Phyllocnistis unipunctella*

Occasional singles taken at light. Distinctive silvery mines can be found in the local Poplars.

*Anthophila fabriciana* – Nettle-tap

Fairly common on the edges of the Golf Course – mostly diurnal. One at light on 6/9.

*Glyphipterix simplicella* – Cocksfoot Moth

A tiny day flying species. Easily found on the Golf Course from mid May-mid June by checking Ox-Eye Daisy flowers, from which they appear to display.

*Argyresthia goedartella*

Singles on 23/7, 15/8 and 19/8.

*Argyresthia curvella*

Singles on 21/6 and 25/6.

*Argyresthia spinosella*

Five on 17/6.

*Argyresthia bonnetella*

Two on 17/6 and 1 on 19/6.

Yponomeuta evonymella – Bird-cherry Ermine  
Singles on 24/7 and 31/7.

Yponomeuta padella – Orchard Ermine  
5/7-3/8. Occasionally 1-3.

Yponomeuta cagnagella – Spindle Ermine  
16/7-24/8. Regularly 1-6.

Pseudoswammerdamia combinella  
One on 29/5.

Swammerdamia compunctella  
One on 24/7.

Paraswammerdamia albicapitella  
Singles on 12/7 and 14/7.

Paraswammerdamia lutarea  
7/7-21/8. Frequently 1-4.

Scythropia crataegella – Hawthorn Moth  
4/7-30/7. 1-2 on four dates with 5 on 24/7.

Ypsolopha dentella – Honeysuckle Moth  
Singles on 23/7, 10/8 and 21/8.

Ypsolopha scabrella  
Singles on 28/7 and 24/8.

Ypsolopha sequella  
7/8-12/9. Singles on five dates.

Plutella xylostella – Diamond-back Moth  
A common immigrant. A total of 3789 from 10/5-3/12. Most frequent in July/August.  
191-319 on six dates 28/7-7/8 with a peak of 1242 on 30/7.

Eidophasia messingiella  
5/6-28/6. Singles on seven dates with 3 on 17/6.

Coleophora alcyonipennella  
One on 24/7.

Coleophora mayrella  
One on 8/7.

Coleophora paripennella  
One on 8/7.

Elachista atricomella  
One on 5/6.

Elaschista argentea  
Easily found resting on the stems of long grass on the golf course from 20/5. Infrequent at light  
from late May-mid June.

Elachista subocellea  
One on 24/7.

- Batia lunaris*  
Singles on 15/7 and 21/7 with 2 on 22/7.
- Batia unitella*  
17/7-7/8. Singles on four dates with 4 on 28/7.
- Hofmannophila pseudospretella* – Brown House-moth  
6/5-31/8. Regularly 1-2 throughout.
- Endrosis sarcitrella* – White-shouldered House-moth  
Two on 16/3. From 10/5-3/9, 1-2 occasionally recorded. Late singles on 3/10 and 12/11.
- Esperia sulphurella*  
A common day flying species on the Golf Course edges from mid May-mid June.
- Carcina quercana*  
19/6-3/9. Common. 1-11 on most nights. Late singles on 23/9, 2/10, 3/10 and 7/10.
- Diurnea fagella*  
One on 20/3.
- Depressaria pastinacella* – Parsnip Moth  
Singles on 29/7 and 8/9.
- Agonopterix heracliiana*  
Two on 21/3 and 1 on 1/4.
- Agonopterix alstroemeriana*  
Following hibernation, singles on 21/3, 28/3 and 29/3. 1-3 on eight dates from 19/7-18/8, after which, a further four singles from 27/9-2/11.
- Agonopterix arenella*  
Following hibernation, 1-3 on four dates from 1/4-19/5. Then singles on 18/9, 28/9 and 11/10.
- Agonopterix nervosa*  
One on 14/9.
- Agonopterix umbellana*  
One on 25/9.
- Ethmia terminella*  
One on 17/6. An uncommon immigrant.
- Metzneria lappella*  
One on 21/6, 2 on 24/7 and 1 on 28/7.
- Metzneria metzneriella*  
One on 19/7 and 2 on 24/7.
- Teleiodes vulgella*  
One on 21/6.
- Bryotropha affinis*  
One on 28/6 and 2 on 23/7.
- Bryotropha terrella*  
21/6-14/8. 1-5 frequently recorded.

*Bryotropha domestica*  
Singles on 1/8 and 7/8.

*Scrobipalpa costella*  
1/10-5/11. Singles on four dates.

*Dichomeris marginella* – Juniper Webber  
Two on 19/6 and singles on nine dates 21/6-5/8.

*Helcystogramma rufescens*  
Singles on 21/6, 19/7 and 26/7.

*Oegoconia quadripuncta*  
Singles on 13/7, 24/7 and 10/8.

*Blastobasis lignea*  
One on 21/6. Then common and noted on most nights from 7/7-4/9.

*Blastobasis decolorella*  
Double brooded. 2/6-31/7. Common, recorded most nights. In smaller numbers and less frequent from 15/9-29/11.

*Mompha raschkiella*  
One on 29/7.

*Limnaecia phragmitella*  
One on 25/7.

*Blastodacna hellerella*  
Singles on 17/6 and 25/6.

*Alucita hexadactyla* – Twenty-plume Moth  
Singles on 10/5 and 15/5. Then 1-2 noted on six dates from 12/7-31/8.

*Agdistis bennetii*  
One on 12/8.

*Marasmarcha lunaedactyla*  
One on 29/7.

*Amblyptilia acanthadactyla*  
Singles on 25/5 and 17/8.

*Platyptilia gonodactyla*  
Singles on 16/6 and 8/8.

*Platyptilia pallidactyla*  
One on 13/7.

*Stenoptilia bipunctidactyla*  
One on 6/6. Singles on 8/8, 13/8, 7/9 and 1/10.

*Stenoptilia pterodactyla*  
One on 7/7.

*Pterophorus pentadactyla* – White Plume Moth  
One on 2/8.

Ovendenia lienigianus

One on 28/6, 2 on 30/7.

Emmelina monodactyla

Following hibernation, noted from 4/3-19/5, regularly 1-3.  
28/8-5/11. Regularly 1-3.

Francis Solly